Ubendol Tab.

(Alibendol 100mg)

COMPOSITION

Each tablet contains Alibendol

·100mg

■ DESCRIPTION

Pink, circle, film-coated tablet

INDICATIONS

Dyspepsia, nausea, vomiting, constipation

■DOSAGE & ADMINISTRATION

300mg~400mg a day, in three to four divided doses, before meals and at bedtime. Dosage may be increased and reduced according to age and symptoms.

■CAUTIONS

- 1. General precautions
 - 1) Consult with doctor or pharmacist before administration to the patients who are administering the other drug.
- When administer for 2 weeks and there is no improvement, stop administration and consult with doctor or pharmacist.
- 2. Children

UBENDOL should not be administered under 7 years old of age.

- 3. Precautions during storage and handling
 - 1) Keep out of reach of children.
 - 2) Store at the place where are avoiding sunlight and if possible, with low humidity and cool and closed tightly.
 - 3) Don't put into other container to prevent misuse and preserve of quality.
- 4. Other
 - 1) Digestive Process: Alibendol stimulates not only the digestion and absorption of fat but also the absorption of fat-soluble vitamines (vit A,D,E,K), calcium and iron by activating the action of Lipase, Glyceridase.
 - 2) Stimulation of cholanopoiesis and biliation: Alibendol show biliation over 3hours continuously. Contrary to hydrocholeretics-secreting low-viscosity blie, Alibendol not only stimulates the secretion of high-concentrated bile but also has longer duration of action than dehydrochloric acid. Also, Alibendol stimulates biliary exretion by relieving abnormal concentration of Oddi's sphincter.
 - 3) Safety: Alibendol can be prescribed safely since the safety is validated with childern (4mg/kg/day) and has no adverse events, contraindications and drug interactions so it can be prescribed.

■STORAGE

Store in tight container at room temperature $(1 \sim 30^{\circ}C)$.

· Keep out of reach of children

Read the instructions before use

